Resonant and non-resonant x-ray scattering studies of mutiferroics

Peter D. Hatton

Department of Physics, Durham University, South Road, Durham, U.K. DH1 3LE Email: p.d.hatton@durham.ac.uk

Multiferroics are a class of strongly correlated electron systems that exhibit more than one primary ferroic order parameter simultaneously in a single phase [1]. Multiferroics have potential applications as actuators, switches, magnetic field sensors and high-density electronic memory devices [2]. In such materials there are strong correlations between the crystalline lattice, charge disproportionation, magnetism and orbital occupancy.

In this talk I will introduce the technique of resonant x-ray scattering in which resonantly enhanced scattering can be observed of superlattice reflections by tuning to appropriate absorption edges. Measurements using polarised x-rays can provide information about both the charge [3] magnetic [4] and orbital ordering [5], as well as minute atomic displacements [6]. More recently, magnetic scattering has been directly observed using non-resonant x-ray scattering [7], which is a more direct probe of magnetism. The talk will conclude with a discussion of how polarised non-resonant soft x-ray scattering can be used to determine the spin and orbital components of the magnetic moment in a complex solid.

References

- [1] S-W. Cheong and M. Mostovoy, Nature Materials 6, 13 (2007).
- [2] R. Ramesh and N.A. Spaldin, Nature Materials 6, 21 (2007).
- [3] E. Nazarenko, J.E. Lorenzo, Y.Joly et al., Phys. Rev. Lett. 97, 056403 (2006).
- [4] R.D. Johnson, S.R. Bland, C. Mazzoli et al., Phys. Rev. B 78, 104407 (2008); T.A.W. Beale, S.B.
- Wilkins, R.D. Johnson et al., Phys. Rev. Lett. 105, 087203 (2010).
- [5] J. Fink, E. Schierle, E. Weschke and J. Geck, Rep. Prog. Phys. 76, 056502 (2013).
- [6] H.C. Walker. F. Fabrizi, L. Paolasini et al., Science 333, 1273 (2011).
- [7] C.Vecchini, A. Bombardi, L.C. Chapon et al., Journal of Physics: Conf. Ser. 519, 012004 (2014).