

Neutron spectroscopy

Andrew Wildes

Institut Laue-Langevin

Plan:

- **Properties of the neutron**
- **Neutron spectroscopy**
- **Harmonic oscillators**
- **Atomic vibrations**
 - Quantized energy levels
 - Tunnelling
- **Magnetic vibrations**
 - Crystal fields
 - Molecular magnets
- **Propagating modes**
 - Phonons
 - Magnons

The neutron

Forms part of the nucleus of the atom

Rest mass (m_n)	$1.675 \times 10^{-27} \text{ kg}$ (c.f. mass of hydrogen = $1.674 \times 10^{-27} \text{ kg}$)
Diameter	$\sim 10^{-15} \text{ m}$ (c.f. diameter of an atom $\sim 10^{-10} \text{ m}$)
Charge	0
Spin	$-1/2$
Magnetic moment	$0.966 \times 10^{-26} \text{ JT}^{-1}$ $-1.913 \mu_N = -\gamma \mu_N$ $1.042 \times 10^{-3} \mu_B$ (c.f. moment on an electron = $1 \mu_B$)

Neutron beams

Through quantum mechanics, neutrons have a *wavelength*.

Energy of a neutron

$$E = h\nu = \hbar\omega$$

h = Planck's constant
 ν = frequency
 ω = angular frequency
 v = speed

$$= \frac{1}{2} m_n v^2$$

Momentum of a neutron

$$p = h/\lambda = \hbar k$$

λ = wavelength
 k = wavenumber

$$\mathbf{p} = \hbar \mathbf{k}$$

‘Standard’ thermal neutrons:

$$v = 2200 \text{ m/s}$$


$$\lambda = 1.798 \text{ \AA}$$

$$T = 293 \text{ K}$$

$$E = 25.3 \text{ meV} = 6 \text{ THz}$$

$$p = 3.68 \times 10^{-24} \text{ kg m s}^{-1} = 3.5 \text{ \AA}^{-1}$$


Neutron scattering – length and time scales



A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Fig. 1.2 Geometry for scattering experiment.



The target volume is initially in state ζ .

A neutron enters with wave vector k and spin s

It interacts with the target.

The final neutron wave vector is k' and spin s' .

The final target state is ζ' .

We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$

$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$

Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n}(k^2 - k'^2)$

Knowing the neutron wavelength

You must know the wavelength to perform a scattering experiment

With neutrons, there are two ways of knowing the wavelength:



1. Use a monochromator
 - Bragg's law: $2d\sin\theta = \lambda$
2. Use time-of-flight
 - Neutron speed $\propto 1/\lambda$, $4\text{\AA} \sim 1000 \text{ m/s}$
 - *chopper* to use *time-of-flight*



– *velocity selector* to monochromate

The neutron spin precession can also give energy change information


Learn to work with vectors



A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Fig. 1.2 Geometry for scattering experiment.



The target volume is initially in state ζ .

A neutron enters with wave vector k and spin s

It interacts with the target.

The final neutron wave vector is k' and spin s' .

The final target state is ζ' .

We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$


$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$

Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n}(k^2 - k'^2)$

A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN



We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$

$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$


Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n} (k^2 - k'^2)$

A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Closing the triangle



We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$

$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$


Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n} (k^2 - k'^2)$

A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Closing the triangle



We measure:

Momentum transfer: $Q = k - k'$

$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$


Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n} (k^2 - k'^2)$

A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Closing the triangle



We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$


$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$

Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n} (k^2 - k'^2)$

A neutron scattering experiment

INSTITUT MAX VON LAUE - PAUL LANGEVIN



We measure:

Momentum transfer: $\mathbf{Q} = \mathbf{k} - \mathbf{k}'$


$$Q^2 = k^2 + k'^2 - 2kk' \cos\Theta$$

Δs

Energy transfer: $\Delta E = \hbar\omega = \frac{\hbar^2}{2m_n}(k^2 - k'^2)$


Classical harmonic oscillators

Pendulum



$$v = \frac{1}{2\pi} \sqrt{\frac{g}{l}}, \quad \omega = \sqrt{\frac{g}{l}}$$

Potential well



$$U = Ax^2$$

$$v = \frac{1}{2\pi} \sqrt{2Ag}, \quad \omega = \sqrt{2Ag}$$

Spring (constant = C)




$$v = \frac{1}{2\pi} \sqrt{\frac{C}{m}}, \quad \omega = \sqrt{\frac{C}{m}}$$

Quantum harmonic oscillators

INSTITUT MAX VON LAUE - PAUL LANGEVIN

$$H\psi = E\psi$$

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + U(x)\psi = E\psi$$






$$U = \frac{1}{2} m\omega^2 x^2$$

$$E = \hbar\omega \left(n + \frac{1}{2} \right)$$

Nitrogen motion in UN

INSTITUT MAX VON LAUE - PAUL LANGEVIN




Endohedral fullerene

NEUTRONS
FOR SOCIETY

“Molecular surgery”


H_2 , D_2 , HD @ C_{60}
 H_2O @ C_{60}

H_2 , $(H_2)_2$ @ C_{70}



K.Komatsu, M.Murata and Y.Murata, Science 307, 238 (2005)

Endohedral fullerene




A. J. Horsewill *et al.*, Proc. Trans. Roy. Soc A **371** (2013) 20110627

C. Beduz *et al.*, PNAS **109** (2012) 12894

Methyl group motion


INSTITUT MAX VON LAUE - PAUL LANGEVIN

Tunnelling in Sodium Acetate Trihydride



Tunnelling in (2,6)lutidine

INSTITUT MAX VON LAUE - PAUL LANGEVIN



With thanks to B. Frick

Magnetism

Magnetism is caused by unpaired electrons or movement of charge.









Magnetic spectroscopy requires a change of neutron *energy*
and *angular momentum* (i.e. the neutron spin changes direction)

Magnetism can be classified as *localized* (i.e. confined to an atomic position)
or *itinerant* (i.e. due to electrons that are moving through the sample)

We'll only discuss localized magnetism today.

Crystal field levels

INSTITUT MAX VON LAUE - PAUL LANGEVIN



Crystal fields in NdPd₂Al₃



INSTITUT MAX VON LAUE - PAUL LANGEVIN

$$H = \sum_{l=2,4,6} B_l^0 O_l^0 + \sum_{l=2,4} B_l^4 O_l^4$$

$O =$ Stevens parameters


(K. W. Stevens, Proc. Phys. Soc A65 (1952) 209)

$B =$ CF parameters,
measured by neutrons



A. Dönni *et al.*, *J. Phys.: Condens. Matter* **9** (1997) 5921


Molecular magnets




Quantum tunneling in Mn₁₂-acetate

INSTITUT MAX VON LAUE - PAUL LANGEVIN


$$H = \sum_{l=2,4,6} B_l^0 O_l^0 + \sum_{l=2,4} B_l^4 O_l^4$$



Calculated energy terms



Neutron spectra



Fitted data with scattering from:

- energy levels
- elastic scattering
- incoherent background

Classical harmonic oscillators

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Spring (constant = C)




$$v = \frac{1}{2\pi} \sqrt{\frac{C}{m}}, \quad \omega = \sqrt{\frac{C}{m}}$$

Propagating lattice vibrations

Take a line of equal masses, M , joined by springs

Take a line of equal masses, M , joined by springs




$$\omega^2 = \frac{4C}{M} \sin^2 \frac{\pi a}{\lambda} = \frac{4C}{M} \sin^2 \frac{ka}{2}$$

Phonon dispersion



INSTITUT MAX VON LAUE - PAUL LANGEVIN

$$\omega^2 = \frac{4C}{M} \sin^2 \frac{\pi a}{\lambda} = \frac{4C}{M} \sin^2 \frac{ka}{2}$$



Phonons in UN

INSTITUT MAX VON LAUE - PAUL LANGEVIN




A.A. Aczel *et al.*, Nat. Comm. **3** (2012) 1124

Phonons in Diamond

INSTITUT MAX VON LAUE - PAUL LANGEVIN




DISPERSION CURVES FOR DIAMOND AT 296 °K



1000
meV


Phonons in fibre DNA

INSTITUT MAX VON LAUE - PAUL LANGEVIN



L. van Eijck *et al.*, PRL 107 (2011) 088102

Phonons in superfluid Helium



B. Fåk *et al.*, PRL **109** (2012) 155305

E. C. Svensson *et al.*, PRB **23** (1981) 4493

D. M. Ceperley and E. L. Pollock, Can. J. Phys. **65** (1987) 1416

H. R. Glyde, *Excitations in Liquid and Solid Helium*
(1994) Clarendon, Oxford

Spin waves and magnons

INSTITUT MAX VON LAUE - PAUL LANGEVIN

A simple Hamiltonian for spin waves is:


$$H = -J \sum_{i,j} \mathbf{s}_i \cdot \mathbf{s}_j$$

J is the magnetic exchange integral, which can be measured with neutrons.


Take a simple ferromagnet:



The spin waves might look like this:




Spin waves have a frequency (ω) and a wavevector (\mathbf{q})



The frequency and wavevector of the waves are *directly measurable* with neutrons

Magnetic excitations in CuSO_4

INSTITUT MAX VON LAUE - PAUL LANGEVIN





Fully polarized state

$$\langle S_n^z \rangle = S$$



Spin wave





Momentum transfer ($h, -1/2, 1/2$)

M. Mourigal *et al.*, Nature Phys. **9** (2013) 435

Spin-waves in $(\text{La,Ba})_2\text{CuO}_4$

INSTITUT MAX VON LAUE - PAUL LANGEVIN





Magnons in FePS₃

INSTITUT MAX VON LAUE - PAUL LANGEVIN



b








Magnons in FePS₃







INSTITUT MAX VON LAUE - PAUL LANGEVIN

$E_i = 31.8 \text{ meV}$



$E_i = 75 \text{ meV}$



[HH0]

D. Lançon *et al.*, PRB **94** (2016) 214407

Conclusions

INSTITUT MAX VON LAUE - PAUL LANGEVIN

Neutron spectroscopy is about measuring quantum oscillations
in solids and liquids

The neutron's momentum and energy are ideal for these

Neutron spectroscopy is a *quantitative* technique that *directly* probes
the dynamics in the sample